

BILINEAR EXPANSIONS OF THE KERNELS OF SOME NONSELFADJOINT INTEGRAL OPERATORS

Milutin Dostanić

Abstract. Let H and S be integral operators on $L^2(0, 1)$ with continuous kernels. Suppose that $H > 0$ and let $A = H(I + S)$. It is shown that if the (nonselfadjoint) operator S is small in a certain sense with respect to H , then the corresponding Fourier series of functions from $R(A)$ (or $R(A^*)$) converges uniformly on $[0, 1]$.

1. Introduction

Let H and S be integral operators on $L^2(0, 1)$ (with inner product $\langle f, g \rangle = \int_0^1 f(x)\overline{g(x)} dx$) with continuous kernels $\mathcal{H}(x, y)$ and $\mathcal{S}(x, y)$ on $[0, 1] \times [0, 1]$. Suppose that $H > 0$ and let

$$A = H(I + S). \quad (1)$$

Classical theorems (case $S = 0$, see [5]) state that the kernel \mathcal{H} can be expanded into a uniformly convergent (on $[0, 1] \times [0, 1]$) bilinear series.

A consequence of this is that every function $f \in R(H)$ has the uniformly convergent Fourier series with respect to the system of eigenfunctions of H . ($R(H)$ denotes the image of H in $L^2(0, 1)$).

Similar results hold in some cases when $S \neq 0$. Namely, if $S = S^*$, it was proved in [1] and [2] that the corresponding variant of Mercer's theorem holds. The proof was based on the spectral theorem for an operator on $L^2(0, 1)$ with the definite or indefinite inner product generated by the formula

$$[f, g] = \langle (I + S)f, g \rangle.$$

In [4], a series of nice results was obtained which were related to bilinear expansions of smooth Carleman's kernels of Mercer type.

A natural question is about bilinear expansions when $S \neq S^*$. We shall show that if the operator S is small in a certain sense with respect to H , then the corresponding Fourier series of functions from $R(A)$ (or $R(A^*)$) converges uniformly

AMS Subject Classification: 47 G 10, 45 P 05

Keywords and phrases: Nonselfadjoint integral operators, bilinear expansion.

on $[0, 1]$. (Note that, because of the continuity of the kernel of A , we have $R(A) \subset C[0, 1]$).

In the sequel, $\mathcal{A}(\cdot, \cdot)$ will denote the kernel of the operator A (defined by (1)).

2. The results

THEOREM. *Let, for the operators H and S from (1), there exists $\omega > 0$ such that $S^*S \leq \omega H^2$. If s_k are singular values of A and f_k the normalized eigenvectors of A^*A (i.e., $A^*A f_k = s_k^2 f_k$) and $g_k = (s_k)^{-1} A f_k$, then the series $\sum_{k \geq 1} s_k \overline{f_k(y)} g_k(x)$ is absolutely convergent on $[0, 1]^2$ and uniformly convergent on $[0, 1]$ with respect to arbitrary variable and its sum is equal to $\mathcal{A}(x, y)$. Also, for every $f \in R(A)$ (resp. $g \in R(A^*)$) the series $\sum_{k \geq 1} \langle f, g_k \rangle g_k$ (resp. $\sum_{k \geq 1} \langle g, f_k \rangle f_k$) converges uniformly on $[0, 1]$ to f (resp. g).*

In the proof of this assertions we need the following two Lemmas.

LEMMA 1. [5] *If $T: L^2(0, 1) \rightarrow L^2(0, 1)$ is the linear operator defined by $Tf(x) = \int_0^1 M(x, y)f(y) dy$ and if $M \in C([0, 1]^2)$ and $\langle Tf, f \rangle \geq 0$ for all $f \in L^2(0, 1)$, then $M(x, x) \geq 0$ for all $x \in [0, 1]$.*

LEMMA 2. *If $A = H(I + S)$, $H > 0$, $S^*S \leq \omega H^2$, then there exists a constant $c > 0$ such that*

$$\sqrt{A^*A} \leq cH, \quad \sqrt{AA^*} \leq cH.$$

Proof. Since

$$A^*A = H^2 + S^*H^2 + H^2S + S^*H^2S \quad (2)$$

we have to estimate $\langle S^*H^2f, f \rangle$ and $\langle H^2Sf, f \rangle$.

The operator H^2 is positive and thus, by the Cauchy inequality, we have

$$\begin{aligned} |\langle S^*H^2f, f \rangle|^2 &= |\langle H^2f, Sf \rangle|^2 \leq \langle H^2f, f \rangle \langle H^2Sf, Sf \rangle \\ &= \|Hf\|^2 \|HSf\|^2 \leq \|Hf\|^2 \|H\|^2 \|Sf\|^2 \\ &= \|Hf\|^2 \|H\|^2 \langle S^*Sf, f \rangle \leq \|Hf\|^2 \|H\|^2 \omega \langle H^2f, f \rangle \\ &= \omega \|H\|^2 \|Hf\|^4. \end{aligned}$$

Therefore

$$|\langle S^*H^2f, f \rangle| \leq \sqrt{\omega} \|H\| \langle H^2f, f \rangle \quad (3)$$

and hence we get

$$|\langle H^2Sf, f \rangle| \leq \sqrt{\omega} \|H\| \langle H^2f, f \rangle. \quad (4)$$

Since

$$\langle S^*H^2Sf, f \rangle = \|HSf\|^2 \leq \|H\|^2 \|Sf\|^2 = \|H\|^2 \langle S^*Sf, f \rangle \leq \omega \|H\|^2 \langle H^2f, f \rangle,$$

from (2), (3), (4) it follows that $\langle A^*Af, f \rangle \leq (1 + \sqrt{\omega} \|H\|)^2 \langle H^2f, f \rangle$, i.e. $A^*A \leq (1 + \sqrt{\omega} \|H\|)^2 H^2$.

Having in mind that the function $\lambda \mapsto \sqrt{\lambda}$ is operator monotone, we get

$$\sqrt{A^*A} \leq (1 + \sqrt{\omega} \|H\|)H. \tag{5}$$

From the equality $A^* = (I + S^*)H$ we get $\|A^*f\| \leq \|I + S^*\| \cdot \|Hf\|$ ($f \in L^2(0, 1)$), i.e.

$$\sqrt{AA^*} \leq \|I + S^*\| \cdot H. \tag{6}$$

From (5) and (6) we obtain the assertion of the Lemma, with

$$c = \max\{1 + \sqrt{\omega} \|H\|, \|I + S^*\|\}. \blacksquare$$

Proof of the Theorem. From $A^*Af_k = s_k^2 f_k$ and $Af_k = s_k g_k$ it follows that $f_k, g_k \in C[0, 1]$ (because A has the continuous kernel) and A has the following singular (see [3]) expansion

$$A = \sum_{k \geq 1} s_k \langle \cdot, f_k \rangle g_k.$$

Also, there holds

$$\begin{aligned} \sqrt{A^*A} f &= \sum_{k \geq 1} s_k \langle f, f_k \rangle g_k \\ \sqrt{AA^*} f &= \sum_{k \geq 1} s_k \langle f, g_k \rangle g_k, \end{aligned} \quad f \in L^2(0, 1). \tag{7}$$

The series on the right-hand side of the previous equalities converge in the norm of $L^2(0, 1)$.

Consider the operators (on $L^2(0, 1)$) S'_n, S''_n defined in the following way: $S'_n = cH - \sum_{k=1}^n \langle \cdot, f_k \rangle f_k, S''_n = cH - \sum_{k=1}^n \langle \cdot, g_k \rangle g_k$. Since, by (7) and Lemma 2, $\langle S'_n f, f \rangle \geq 0, \langle S''_n f, f \rangle \geq 0, f \in L^2(0, 1)$ and since the operators S'_n, S''_n have continuous kernels, we get from Lemma 1

$$c\mathcal{H}(x, x) \geq \sum_{k=1}^n s_k |f_k(x)|^2, \quad c\mathcal{H}(x, x) \geq \sum_{k=1}^n s_k |g_k(x)|^2, \quad n \in \mathbf{N}.$$

Since $\mathcal{H} \in C([0, 1]^2)$, there exists $M_0 < +\infty$ such that

$$\sum_{k=1}^n s_k |f_k(x)|^2 \leq M_0, \quad \sum_{k=1}^n s_k |g_k(x)|^2 \leq M_0. \tag{8}$$

From (8) it follows that the series

$$\sum_{k \geq 1} s_k \overline{f_k(y)} g_k(x)$$

is absolutely convergent for all $x, y \in [0, 1]$. Let $S(x, y)$ denote its sum. Observe that from (8) it follows that the partial sums of the previous series are bounded by M_0 .

Fix $x \in [0, 1]$. Then we have

$$\begin{aligned} \left| \sum_{k=p}^q s_k \overline{f_k(y)} g_k(x) \right|^2 &\leq \sum_{k=p}^q s_k |f_k(y)|^2 \sum_{k=p}^q s_k |g_k(x)|^2 \\ &\leq M_0 \sum_{k=p}^q s_k |g_k(x)|^2 \rightarrow 0 \quad (p, q \rightarrow \infty) \end{aligned}$$

and the series $\sum_{k \geq 1} s_k \overline{f_k(y)} g_k(x)$ converges uniformly with respect to y on $[0, 1]$, for every fixed x and hence its sum is a continuous function with respect to y .

Let $f \in C[0, 1]$ be a fixed function. Then (because of the uniform convergence with respect to y)

$$\int_0^1 S(x, y) f(y) dy = \sum_{k \geq 1} s_k g_k(x) \int_0^1 f(y) \overline{f_k(y)} dy = \sum_{k \geq 1} s_k g_k(x) \langle f, f_k \rangle. \quad (9)$$

(The series on the right-hand side of (9) converges not only for every x but also uniformly with respect to x because

$$\begin{aligned} \left| \sum_{k=p}^q s_k g_k(x) \langle f, f_k \rangle \right|^2 &\leq \sum_{k=p}^q s_k |g_k(x)|^2 \sum_{k=p}^q s_k |\langle f, f_k \rangle|^2 \\ &\leq M_0 s_p \|f\|^2 \rightarrow 0 \quad (p, q \rightarrow \infty) \end{aligned}$$

On the other hand, from the singular expansion of A we get

$$\int_0^1 \mathcal{A}(x, y) f(y) dy = \sum_{k \geq 1} s_k g_k(x) \langle f, f_k \rangle \quad (10)$$

(the series converges in the norm of $L^2(0, 1)$).

Thus, from (9), (10) it follows that for every $f \in C[0, 1]$ we have

$$\int_0^1 (\mathcal{A}(x, y) - S(x, y)) f(y) dy = 0.$$

Putting $f(y) = \overline{S(x, y)} - \overline{\mathcal{A}(x, y)}$ ($\in C[0, 1]$) we get

$$\int_0^1 |\mathcal{A}(x, y) - S(x, y)|^2 dy = 0$$

and hence $\mathcal{A}(x, y) = S(x, y)$ for every $y \in [0, 1]$. Since $x \in [0, 1]$ was arbitrary, we have $\mathcal{A}(x, y) = S(x, y)$, $x, y \in [0, 1]$. So

$$\mathcal{A}(x, y) = \sum_{k \geq 1} s_k \overline{f_k(y)} g_k(x)$$

for every $x, y \in [0, 1]$.

Let now $f \in R(A)$. Then

$$f(x) = \int_0^1 \mathcal{A}(x, y) \varphi(y) dy, \quad \varphi \in L^2(0, 1)$$

and thus, by the Lebesgue dominated convergence theorem, we have ($\mathcal{A}_n(x, y) = \sum_{k=1}^n s_k \overline{f_k(y)} g_k(x)$)

$$\begin{aligned} f(x) &= \int_0^1 \lim_{n \rightarrow \infty} \mathcal{A}_n(x, y) \varphi(y) dy = \lim_{n \rightarrow \infty} \int_0^1 \mathcal{A}_n(x, y) \varphi(y) dy \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^n s_k g_k(x) \langle \varphi, f_k \rangle = \sum_{k \geq 1} s_k g_k(x) \langle \varphi, f_k \rangle. \end{aligned}$$

Since the preceding series converges uniformly with the respect to $x \in [0, 1]$ and since $\{g_k\}$ is an orthonormal system in $L^2(0, 1)$ (see [3]), we have $s_k \langle \varphi, f_k \rangle = \langle f, g_k \rangle$ and, finally, we get

$$f(x) = \sum_{k \geq 1} \langle f, g_k \rangle g_k(x)$$

(the series converges uniformly on $[0, 1]$).

The assertion for the function $g \in R(A^*)$ can be proved in a similar way. ■

REMARK. The second part of the Theorem was proved in [5] in a different way. The proof presented here is a consequence of the previously established bilinear expansion of the function $\mathcal{A}(x, y)$.

REFERENCES

- [1] M. R. Dostanić, *Generalizations of Mercer's theorem for a class of nonselfadjoint operators*, Mat. Vesnik **41** (1989), 22–29.
- [2] M. R. Dostanić, *Generalizations of the Mercer theorem*, Publ. de l'Inst. Math., Nouv. serie, **54(68)** (1993), 63–70.
- [3] I. C. Gohberg and M. G. Krein, *Introduction to the Theory of Linear Nonselfadjoint Operators*, Translation of the Math. Monographs, **18**, Amer. Math. Soc., Providence, R.I., 1969.
- [4] I. M. Novitskii, *Integral representations of linear operators by smooth Carleman kernels of Mercer type*, Proc. London Math. Soc. (3) **68** (1994), 161–177.
- [5] F. Riesz, B. Sz.-Nagy, *Leçons d'analyse fonctionnelle*, Akademiai Kiado, Budapest, 1972.

(received 05.05.2001, in revised form 10.04.2002)

Faculty of Mathematics, University of Belgrade, Studentski trg 16, 11000 Beograd, Yugoslavia
E-mail: domi@matf.bg.ac.yu